

Quantum computing and communication with decoherence-free atomic states

Mang Feng *

Laboratory of Magnetic Resonance and Atomic and Molecular Physics,

Wuhan Institute of Physics and Mathematics, Academia Sinica, Wuhan, 430071, P.R.China

(February 1, 2008)

Abstract

To resist decoherence from destroying the phase factor of qubit state, it is important to use decoherence-free states for processing, transmitting and storing quantum information in quantum computing and quantum communication. We propose a practical scheme using four atoms with decoherence-free states in a single-mode cavity to realize the entanglement and fundamental quantum logic gates. The transmission of quantum information can be made directly from one atom to another, in which the cavity is only virtually excited. The possible application and the experimental requirement of our proposal are discussed.

PACS numbers: 03.67.Lx, 03.65.Bz

Typeset using REVTeX

*Present address: Institute for Scientific Interchange Foundation, Villa Gualino, Viale Settimio Severo 65, I-10131, Torino, Italy. Electronic address: feng@isiosf.isi.it

Both quantum computing and quantum communication have been drawn much attention over past few years. It is predicted that quantum computing would outperform the computation made by existing computers in treating some problems, such as solving classically intractable problems [1,2] and finding tractable solutions more rapidly [3]. There are many schemes for realizing quantum computing with nuclear magnetic resonance (NMR) technique [4], trapped ions [5,6], cavity quantum electrodynamics (QED) [7], and so on. It has been proven [8] that a suitable combination of some single-qubit rotations and two-qubit operations will in principle construct any quantum computing operation we desired. Up to now, some demonstrative experiments have been made to show the possibility of quantum computing in NMR [9], ion trap [10,11] and cavity-atom system [12].

Besides the application in quantum computing, cavity QED technique is more suitable for quantum communication [13,14]. To our knowledge, there are some proposals for teleportation by means of conditional dynamics in cavity-atom systems [15]. Both for the microwave cavity and optical one, as long as the cavity with high-quality factor is initially cooled down to the vacuum state, quantum information will be transferred from one atom to the cavity, and then from the cavity to another atom. Sometimes the cavity decay is also useful for teleportation of an atomic state [16]. Recently, a novel scheme [17] was proposed to use cavities in thermal states for teleportation of an atomic state, in which two identical atoms are required to be put into the cavity simultaneously.

Quantum state is too fragile to exist for a long time unless the system under consideration is perfectly isolated from its environment. To overcome the detrimental effect of decoherence, some methods have been proposed for correcting errors of quantum states after mistakes have been detected [18], or employing the suitable dynamical coupling to average out the detrimental effect from the environment [19]. Alternatively, we can also use some suitable encoded states to resist the decoherence effect [20]. As far as we know, some studies of logic gates on qubits encoded with decoherence-free states have been made before in the context of cavity QED [21] and solid state system [22]. In recent experiments with photons [23] and trapped ions [24], the decoherence-free states, which prevent decoherence from destroying

the phase factors in quantum states of qubits, have been demonstrated to be robust against the collective dephasing. The main idea of these experiments is to encode the two-particle quantum state to be $(|eg\rangle + e^{i\theta}|ge\rangle)/\sqrt{2}$ with $|e\rangle$ and $|g\rangle$ being excited and ground states respectively of each particle, and θ the phase factor, so that no phase change would happen under collective dephasing. In fact, this idea was also used in a recent work [25] for Grover search with pairs of trapped ions, in which such a decoherence-free state can also eliminate the change of phase factor due to the time evolution of different components of quantum superposition.

In this contribution, we would demonstrate the implementation of entanglement and fundamental quantum logic gates with four-particle decoherence-free states in the cavity-atom system, if four identical atoms can be put into a single-mode cavity simultaneously. We would show that the entanglement of four atoms can be made in one step, much more quickly than any other methods proposed previously. More importantly, the realization of fundamental quantum logic gates would open a new way to quantum computing and quantum communication with decoherence-free state in the context of cavity QED. Finally we will discuss the application of our scheme by taking an example of teleportation.

Consider four identical two-level atoms experiencing simultaneously a monochromatic cavity field, with the Hamiltonian

$$H = H_0 + H_{int} \quad (1)$$

where $H_0 = \omega_a \sum_{i=1}^4 \sigma_{iz} + \omega a^\dagger a$ and $H_{int} = G \sum_{i \neq j; i,j=1}^4 (a^2 \sigma_i^+ \sigma_j^+ + a^{+2} \sigma_i \sigma_j)$ with ω_a and ω being respectively resonant transition frequency of two levels of each atom and the frequency of the cavity field. $a^\dagger$ and a are creation and annihilation operators of the cavity mode respectively. G is the coupling strength between the cavity and the atom, and σ_i ($i = +, -$ and z) are familiar Pauli operators. In the case of large detuning between ω_a and ω , there is no energy exchange between atoms and the cavity. So the possible transitions in this system are between $|egeg\rangle$ and $|gege\rangle$, or between $|geeg\rangle$ and $|egge\rangle$, or between $|eegg\rangle$ and $|ggee\rangle$ with $|e\rangle$ and $|g\rangle$ being excited and ground states of each atom

and $|n\rangle$ the cavity state. The Rabi frequency Ω of these transitions, via intermediate state $|gggg\ n+2\rangle$ and $|eeee\ n-2\rangle$, can be calculated by second order perturbation theory. For example, for the transition between $|egeg\ n\rangle$ and $|gege\ n\rangle$, we have

$$\Omega = \frac{\langle egeg\ n|H_{int}|gggg\ n+2\rangle\langle gggg\ n+2|H_{int}|gege\ n\rangle}{\delta} + \frac{\langle egeg\ n|H_{int}|eeee\ n-2\rangle\langle eeee\ n-2|H_{int}|gege\ n\rangle}{-\delta} = (4n+2)G^2/\delta. \quad (2)$$

For other two cases, we can obtain the same value of Ω as in Eq.(2). Different from the work of two particles [6,17], the Rabi frequency in the present system is related to the cavity state. For simplicity, we first assume the cavity state to be initially the vacuum state. But this requirement will be loosened, as discussed later. When $n=0$, the effective Hamiltonian of Eq.(1) is $H^{eff} = H_0^{eff} + H_{int}^{eff}$ with $H_0^{eff} = \Omega \sum_{i=1}^4 |e_i\rangle\langle e_i| a^2 a^{+2}$ and

$$H_{int}^{eff} = \Omega(\sigma_1^+ \sigma_2^+ \sigma_3^- \sigma_4^- + \sigma_1^+ \sigma_2^- \sigma_3^+ \sigma_4^- + \sigma_1^+ \sigma_2^- \sigma_3^- \sigma_4^+ + \sigma_1^- \sigma_2^+ \sigma_3^+ \sigma_4^- + \sigma_1^- \sigma_2^+ \sigma_3^- \sigma_4^+ + \sigma_1^- \sigma_2^- \sigma_3^+ \sigma_4^+) \quad (3)$$

where Pauli operators are written as $\sigma_i^\pm = I \otimes \dots \otimes \sigma^{(i)\pm} \otimes \dots \otimes I$ with $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $\sigma^{(i)\pm}$ denoting that the i th matrix is σ^+ or σ^- . For example, $\sigma_1^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $\sigma_2^- = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

As H_0^{eff} describes Stark shifts related to the photon number, which would become a phase factor in the wave function, in what following, we would only focus on studying the term of H_{int}^{eff} . By means of Eq.(3), the straightforward deduction for the time evolution of the system in the interaction representation is

$$\begin{aligned} |egeg\rangle &\rightarrow \cos(\Omega t)|egeg\rangle - i\sin(\Omega t)|gege\rangle, |gege\rangle \rightarrow \cos(\Omega t)|gege\rangle - i\sin(\Omega t)|egeg\rangle, \\ |egge\rangle &\rightarrow \cos(\Omega t)|egge\rangle - i\sin(\Omega t)|geeg\rangle, |geeg\rangle \rightarrow \cos(\Omega t)|geeg\rangle - i\sin(\Omega t)|egge\rangle, \\ |eegg\rangle &\rightarrow \cos(\Omega t)|eegg\rangle - i\sin(\Omega t)|ggee\rangle, |ggee\rangle \rightarrow \cos(\Omega t)|ggee\rangle - i\sin(\Omega t)|eegg\rangle \end{aligned} \quad (4)$$

which means that we can obtain the four-atom entangled state in one step as long as the four atoms were put into the cavity simultaneously. By setting different initial conditions of each atoms and controlling the evolution time, we would have different four-atom entangled states. As the entanglement of four atoms in our scheme can be made much more quickly than any other known methods, our scheme would be useful in quantum communication for preventing decoherence from affecting quantum channels.

The four-particle entanglement presented here is different from the entanglement of four trapped ions achieved in NIST [11]. The form of the four-ion entangled state is $\Phi_{en} = \frac{1}{\sqrt{2}}(|1111\rangle + |0000\rangle)$ with $|1\rangle$ and $|0\rangle$ being the excited and ground states respectively of each ion because the trapped ions exchange energy with laser beams without changing vibrational states of the ions, and finally get to Φ_{en} . In contrast, in our scheme, as there is no laser beam in the system, atoms only exchange energy with each other in the case of no change of cavity state. Therefore we obtain different forms of four-particle entangled states. Another difference of our scheme from Ref.[11] is that, the coupling between atoms and cavity in our scheme is dependent on the cavity state although the cavity state is only virtually excited in atomic state transitions. In the treatment above, we supposed the cavity state to be initially vacuum. While for the cavity state with $n = 1, 2, \dots$, our scheme could be still available. However, if n is too large, the perturbation theory will be invalid for solving the coupling Ω . In this sense, even if the cavity state is in thermal state, our proposal would still work as long as the mean photon number in the cavity is small.

Similar to Ref.[6] that the controlled-NOT (CNOT) gate has been constructed on two identical ions experiencing a single-mode laser beam, we can also carry out the CNOT gate on the four identical atoms based on Eq.(4). We choose four states $|egeg\rangle_{1234}$, $|gege\rangle_{1234}$, $|egge\rangle_{1234}$ and $|geeg\rangle_{1234}$ to form the computational space where subscripts mean the labeling of the atoms. Following sequence of operation: $H_{34}, P_{34}, R, P_{34}, H_{12}, H_{34}$ and P_{34}^{-1} will yield the CNOT gate as $|eg\rangle_{12} |ge\rangle_{34} \rightarrow |eg\rangle_{12} |ge\rangle_{34}$, $|ge\rangle_{12} |ge\rangle_{34} \rightarrow |ge\rangle_{12} |ge\rangle_{34}$, $|eg\rangle_{12} |eg\rangle_{34} \rightarrow |ge\rangle_{12} |eg\rangle_{34}$ and $|ge\rangle_{12} |eg\rangle_{34} \rightarrow |eg\rangle_{12} |eg\rangle_{34}$, where H_{ij} is made by putting the atoms of i and j into a single-mode cavity simultaneously with the

transformations $|eg\rangle_{ij} \rightarrow (|eg\rangle_{ij} - i|ge\rangle_{ij})/\sqrt{2}$ and $|ge\rangle_{ij} \rightarrow (|ge\rangle_{ij} - i|eg\rangle_{ij})/\sqrt{2}$ [17]. P_{ij} is a $\frac{\pi}{2}$ phase change of $|eg\rangle_{ij}$ and R is the evolution shown in Eqs.(4) with $\Omega t = 3\pi/4$.

As the Hadamard transformation on two atoms can be realized by H_{ij} in addition to P_{ij} , the realization of CNOT gates means that any quantum computing operation can be constructed with decoherence-free states in cavity-atom systems. Meanwhile, our scheme would also have important application in teleportation. In the original proposal of teleportation, an unknown quantum state could be perfectly sent from Alice to Bob over arbitrary distance by a measurement of Alice and suitable operations of Bob, in which an entangled state shared previously by Alice and Bob and two bits of classical message sent from Alice to Bob are necessary. However, if we want to teleport an atomic superposition state with components of different eigenenergies, the receiver would probably obtain another state instead of the desired one, because of the time evolution of component states. For example, Alice wants to send the state of $\psi = |g\rangle + e^{i\theta}|e\rangle$ to Bob, where $|g\rangle$ and $|e\rangle$ are different eigenstates respectively. According to Schrödinger equation, during a time interval t , ψ will evolve to $e^{-iE_g t/\hbar}|g\rangle + e^{i\theta}e^{-iE_e t/\hbar}|e\rangle$ with E_g and E_e being the eigenenergies of states $|g\rangle$ and $|e\rangle$ respectively. So if Alice takes a period of T to inform Bob about her measurement result, Bob will obtain $|g\rangle + e^{i\theta}e^{-i(E_e - E_g)T/\hbar}|e\rangle$. It means that Bob could not obtain the desired state ψ unless teleportation is made superluminally (i.e., $T = 0$) or in the system with $E_g = E_e$. This evolution problem of eigenstates exists in all teleportation schemes with cavity QED [15-17]. But no consideration has yet been paid on this respect. While if teleported state is set to be $\psi = |ge\rangle + e^{i\theta}|eg\rangle$, and Bell states are $\Phi^\pm = (|egeg\rangle \pm i|gege\rangle)/\sqrt{2}$ and $\Psi^\pm = (|egge\rangle \pm i|geeg\rangle)/\sqrt{2}$, by supposing $|eg\rangle = |\tilde{1}\rangle$ and $|ge\rangle = |\tilde{0}\rangle$, we can perfectly realize the teleportation idea in Ref.[13] with cavity QED technique. We should emphasize that Bell states can be fully distinguished in our scheme, instead of only $\Psi^\pm$ being identified as in Ref.[17]. If we put the four atoms with Bell states into a single-mode cavity for a period of $t = \pi/4\Omega$, we have the time evolution yielding $\Phi^+ \rightarrow |egeg\rangle$, $\Phi^- \rightarrow -i|gege\rangle$, $\Psi^+ \rightarrow |egge\rangle$ and $\Psi^- \rightarrow -i|geeg\rangle$. So the individual measurement on the atoms would

make us identify each of Bell states.

Let us discuss the possible error happening in the implementation of the CNOT gate. Our scheme for CNOT gate includes both two-atom and four-atom operations. Any inadequate operation on two-atom pairs (with H_{ij} and P_{ij}) or four atoms (with R) would produce errors in the CNOT gate. For convenience of our discussion, we consider a simple case. Suppose when we put four atoms with $|egeg\rangle_{1234}$ into the cavity, an imperfect manipulation takes place, i.e., two atoms 1 and 2 being put into the cavity a little sooner than the atoms 3 and 4, which yields $\Psi = \cos(\lambda t_1)\{\cos[\Omega(t - t_1)]|egeg\rangle_{1234} - i\sin[\Omega(t - t_1)]|gege\rangle_{1234}\} - i\sin(\lambda t_1)\{\cos[\Omega(t - t_1)]|geeg\rangle_{1234} - i\sin[\Omega(t - t_1)]|egge\rangle_{1234}\}$ where $\lambda = \Omega/2$ [17], t_1 is the time for atoms 1 and 2 in the cavity before the atoms 3 and 4 are input, and we neglect the common phase factor. If we hope to have a operation R , we will find that the fidelity is $\langle \Psi_R | \Psi \rangle \approx 0.98$ in the case of $t_1 = 0.02t$ with Ψ_R the desired state under the operation R . So our proposal still works very well [26]. Moreover, how to distinguish the four identical atoms is essential to our scheme. Above discussion has shown that any confusion in the manipulation of atoms will make our proposal incorrect or even invalid. Fortunately, four identical atoms in a cavity are distinguishable with the present cavity QED technique, because the atoms in the cavity can be manipulated at centimeter-scale distance [27]. Furthermore, the spontaneous emission of the excited state is another possible detrimental effect in our scheme. From the discussion above, we know that the time for achieving a maximal entanglement and a CNOT gate of four atoms are respectively $\pi\delta/8G^2$ and $7\pi\delta/8G^2$. As $G = 2\pi \times 47kHz$ and the lifetime of the excited state of a Rydberg atom is generally $3 \times 10^{-2}s$ [27], if $\delta = 10G$, the time of above implementations would be on the order of $10^{-4}s$, which is much shorter than the lifetime of the excited state of a Rydberg atom. So our scheme is practical.

To summarize, a practical proposal for quantum communication and fundamental gates of quantum computing with four-atom decoherence-free states has been put forward, based on the supposition that we can simultaneously put four identical atoms into a single-mode cavity with few photons. As the entanglement and quantum gates are made between atoms,

instead of between atoms and photons, the cavity state is only virtually excited so that the cavity is not strictly required to be initially in vacuum state and the quality factor of the cavity is not necessarily set to be very high [17]. Moreover, the information stored in atoms with decoherence-free states is more safe than that stored in photons, and the implementation on atoms for processing, transmitting or storing quantum information is easier than that on photons. Furthermore, the former works involving photons had no way to completely measure Bell states due to the lack of photon-photon interaction [28]. While in our scheme, the effective interaction in the cavity-atom system guarantees the complete measurement of Bell states. Besides although the four-particle entanglement has been realized previously in the ion trap, the four-particle entanglement proposed here is different and is more important because it corresponds to decoherence-free states. While except the dephasing, we have not considered any other effects of decoherence in our scheme, because the collective dephasing is much stronger than other effects of decoherence [23,24] when the atoms are closed enough. Compared with previous proposals [7,17], our scheme is not more stringent. It can be experimentally achieved with the cavity QED technique presently or in the near future. We hope that our proposal would be useful for perfect information processing, transmission and storage in quantum computing and quantum communication.

Valuable discussion with Jianwei Pan is highly acknowledged. The author also would like to thank the hospitality of Max Planck institute for the Physics of Complex Systems, where part of this work was carried out. The work is partly supported by the National Natural Science Foundation of China.

REFERENCES

- [1] P.W. Shor, in the Proceeding of the 35th Annual symposium on Foundation of Computer Science (IEEE Computer Society Press, Los Alamitos, CA, 1994), P. 116; A. Ekert and R. Jozsa, *Rev.Mod.Phys.* **68**, 733 (1996).
- [2] T. Hogg, *Phys. Rev. Lett.* **80**, 2473 (1998).
- [3] L.K. Grover, *Phys.Rev.Lett.* **79**, 325 (1997); *ibid*, **79**, 4709 (1997); *ibid*, **80**, 4329 (1998).
- [4] D.P. DiVincenzo, *Phys.Rev.A* **52**, 1015 (1995); H. Schack and C.M. Caves, *Phys.Rev.A* **60**, 4354 (1999).
- [5] J.I. Cirac and P. Zoller, *Phys.Rev.Lett.* **74**, 4091 (1995); C. Monroe *etal*, *Phys.Rev.A* **55**, R2489 (1997); J.F. Poyatos, J.I. Cirac, and P. Zoller, *Phys.Rev.Lett.* **81**, 1322 (1998); J.I. Cirac, and P. Zoller, *Nature*, **404**, 579 (2000); L.M. Duan, J.I. Cirac, and P. Zoller, *Science*, **292**, 1695 (2001).
- [6] K. Mølmer and A. Sørensen, *Phys.Rev.Lett.* **82**, 1935 (1999); A. Sørensen and K. Mølmer, *Phys.Rev.Lett.* **82**, 1971 (1999); *Phys.Rev.A* **62**, 022311 (2000).
- [7] A. Barenco, D. Deutsch and A. Ekert, *Phys.Rev.Lett.* **74**, 4083 (1995); P. Domokos *etal*, *Phys.Rev.A* **52**, 3554 (1995).
- [8] T. Sleator and H. Weinfurter, *Phys.Rev.Lett.* **74**, 4087 (1995); S. Lloyd, *Phys.Rev.Lett.* **75**, 346 (1995); D.P. DiVincenzo, *Phys.Rev.A* **51**, 1015 (1995); A. Barenco *et al*, *Phys.Rev.A* **52**, 3457 (1995).
- [9] I.L. Chuang, N. Gershemfeld and M. Kubinec, *Phys.Rev.Lett.* **80**, 3408 (1998); I.L. Chuang *et al*, *Nature*, **393**, 393 (1998); D.G. Cory *et al*, *Phys.Rev.Lett.* **81**, 2152 (1998); D.G. Cory, M.D. Price and T.F. Havel, *Physica D* **120**, 82 (1998); E. Knill, R. Laflamme, R. Martinez and C.-H. Tseng, *Nature*, **404**, 368 (2000).
- [10] C. Monroe *et al*, *Phys.Rev.Lett.* **75**, 4714 (1995); Q.A. Turchett *et al*, *Phys.Rev.Lett.*

- 81**, 3631 (1998); H.C. Nägerl *et al*, Phys.Rev.A **60**, 145 (1999); M.A. Rowe *et al*, Nature **409**, 791 (2001).
- [11] C.A. Sackett *et al*, Nature, **404**, 256 (2000).
- [12] Q.A. Turchette *et al*, Phys.Rev.Lett. **75**, 4710 (1995); P.W.H. Pinkse, T. Fischer, P. Maunz and G. Rempe, Nature **404**, 365 (2000).
- [13] C.H. Bennett *etal*, Phys.Rev. Lett. **70**, 1895 (1993).
- [14] C.H. Bennett and S.J. Wiesner, Phys.Rev.Lett. **69**, 2882 (1992).
- [15] L. Davidovich *et al*, Phys.Rev.A **50**, R895 (1994); J.I. Cirac and A.S. Parkins, Phys.Rev.A **50**, R4441 (1994); M.H.Y. Moussa, Phys.Rev.A **55**, R3287 (1997).
- [16] S. Bose, P.L. Knight, M.B. Plenio and V. Vedral, Phys.Rev.Lett, **83**, 5158 (1999).
- [17] S.B. Zheng and G.C. Guo, Phys.Rev.Lett. **85**, 2392 (2000); S.B. Zheng and G.C. Guo, Phys.Rev.A **63**, 044302 (2001).
- [18] D.Gottesman, Phys.Rev.A **54**, 1862 (1996); C.H.Bennett and P.W.Shor, IEEE Trans.Inf.Theory **44**, 2724 (1998).
- [19] L. Viola, E.Knill and S.Lloyd, Phys.Rev.Lett. **82**, 2417 (1999).
- [20] L.M. Duan and G.C. Guo, Phys.Rev.Lett. **79**, 1953 (1997); P. Zanardi and M. Rasetti, Phys.Rev.Lett. **79**, 3306 (1997); D.A. Lidar, D. Bacon, and K.B. Whaley, Phys.Rev.Lett. **82**, 4556 (1999).
- [21] A. Beige, D. Braun, B. Tregenna and P.L. Knight, Phys.Rev.Lett. **85**, 1762 (2000).
- [22] D. Bacon, J. Kempe, D.A. Lidar, and K.B. Whaley, Phys.Rev.Lett. **85**, 1758 (2000).
- [23] P.G.Kwiat, A.J.Berglund, J.B.Altepeter and A.G.White, Science **290**, 498 (2000).
- [24] D. Kielpinski *et al*, Science **291**, 1013 (2001).

- [25] M. Feng, Phys.Rev.A **63**, 052308 (2001).
- [26] For other cases, such as one atom is input before other three and so on, the calculation is a little complicated. But as long as the time t_1 is small enough, for example, less than $0.05t$, the fidelity would be still very high.
- [27] A. Rauschenbeutel *et al*, Science **288**, 2024 (2000).
- [28] D. Bouwmeester *et al*, Nature **390**, 575 (1997); S.L. Braunstein, A. Mann, M. Revzen, Phys.Rev.Lett. **68**, 3259 (1992).